Manage postpartum hemorrhage.

Bakri

POSTPARTUM BALLOON WITH RAPID INSTILLATION COMPONENTS




A simple solution for postpartum hemorrhage (PPH)

Sequential treatment considerations for conservative management of uterotonic-resistant PPH¹

Vaginal birth

- 1 Bimanual massage
- 2 Repair of any lacerations
- 3 Dilation and curettage to rule out retained placenta
- 4 Placement of intrauterine balloon²




5 Selective embolization

Cesarean birth

- 1 Fundal massage
- Inspection of the broad ligament and posterior uterus and inspection for the presence of retained placenta
- Placement of intrauterine balloon²





4 Uterine artery ligation

Note: The above flowchart describes sequential treatment considerations for uterotonic-resistant PPH when blood loss is less than 1,500 mL and after evaluation for uterine inversion and amniotic fluid embolism.

^{1.} California Maternal Quality Care Collaborative. Obstetric hemorrhage emergency management plan: table chart format. Version 2.0. Published 2014. Accessed August 31, 2016.

^{2.} Please refer to the IFU for detailed indications, contraindications, warnings, and precautions.






Bakri^{*}

POSTPARTUM BALLOON WITH RAPID INSTILLATION COMPONENTS

This device is intended to provide temporary control or reduction of postpartum uterine bleeding when conservative management is warranted. The set includes a syringe, 180 cm long extension tubing with bag spike, and dual check valve. Please refer to the Instructions for Use for complete information on product usage, indications, and contraindications.



Order	Reference	Catheter	Length	Volume
Number	Part Number	Fr	cm	mL
G24237	J-SOSR-100500	24	54	500

Some products or part numbers may not be available in all markets. Contact your local Cook representative or Customer Service for details.



Scan this OR code to see other resources.

Resources

- of a positive 'tamponade test' in the management of postpartum haemorrhage. BJOG. 2010;117(3):295-303.
- 2. Vintejoux E, Ulrich D, Mousty E, et al. Success factors for Bakri $^{\text{TM}}$ balloon usage secondary to uterine atony: a retrospective, multicentre study. Aust NZJ Obstet Gynaecol. 2015;55(6):572-577.

1. Georgiou C. Intraluminal pressure readings during the establishment 3. Vitthala S, Tsoumpou I, Anjum ZK, et al. Use of Bakri balloon in postpartum haemorrhage: a series of 15 cases. Aust NZJ Obstet Gynaecol. 2009;49(2):191-194.

Customer Service

EU Website: cookmedical.eu EDI: cookmedical.eu/edi

Distributors: +353 61239240, ssc.distributors@cookmedical.com

Austria: +43 179567121, oe.orders@cookmedical.com Belgium: +32 27001702, be.orders@cookmedical.com Denmark: +45 38487607, da.orders@cookmedical.com Finland: +358 972519996, fi.orders@cookmedical.com France: +33 171230269, fr.orders@cookmedical.com Germany: +49 6950072804, de.orders@cookmedical.com Hungary: +36 17779199, hu.orders@cookmedical.com Iceland: +354 800 7615, is.orders@cookmedical.com Ireland: +353 61239252, ie.orders@cookmedical.com Italv: +39 0269682853, it.orders@cookmedical.com Netherlands: +31 202013367, nl.orders@cookmedical.com

Norway: +47 23162968, no.orders@cookmedical.com

Spain: +34 912702691, es.orders@cookmedical.com Sweden: +46 858769468, se.orders@cookmedical.com Switzerland - French: +41 448009609, fr.orders@cookmedical.com Switzerland - Italian: +41 448009609, it.orders@cookmedical.com Switzerland - German: +41 448009609, de.orders@cookmedical.com United Kingdom: +44 2073654183, uk.orders@cookmedical.com

USA Website: cookmedical.com EDI: cookmedical.com/edi.do

Americas:

Phone: +1 812.339.2235, 800.457.4500, Fax: 800.554.8335 E-mail: customersupport@cookmedical.com

Australia:

Phone: +61 734346000, 1800777222, Fax: +61 734346001, 1800077283

E-mail: cau.custserv@cookmedical.com



